## IN THE CLAIMS

- 1. (Currently Amended) A method of conducting R chemical reactions, where R is a positive integer> 1, in a system which includes an apparatus which provides energy for the chemical reactions, said system also including a parameter selecting unit having a user interface and storage means for carrying a database, said chemical reaction involving one or more chemical species  $^{X}B$  and resulting in a reaction product  $^{X}D$  which includes a functionality  $\delta$ , where the chemical reaction involves one or more functionalities  $\beta$  in the  $^{X}B$ 's which are transformed into  $\delta$  in  $^{X}D$ , each reaction being performed under the influence of one or more corresponding chemical substances  $A_{R}$ , such chemical substances  $A_{R}$  including a chemical functionality  $\alpha_{R}$  being involved in the transformation of the functionality/functionalities  $\beta$  to the functionality  $\delta$ , said database comprising N sets of associated data, each of the N sets comprising
  - i) a set of reaction parameters for a chemical reaction involving the transformation of one or more functionalities  ${}^{N}\beta$  of chemical species  ${}^{N}B$  into  ${}^{N}\delta$  in a product  ${}^{N}D$  under the influence of one or more chemical substances  ${}^{N}A$ , such chemical substance(s) each including a chemical functionality  ${}^{N}\alpha$  being involved in the transformation of the functionality  ${}^{N}\beta$  to the functionality  ${}^{N}\delta$ ; and
- ii) functional or structural information about the chemical species <sup>N</sup>B; the method comprising that
- \* the user provides information to the user interface of the parameter selection unit about the functionality/functionalities  $\beta$  in the chemical species  $^{X}B$ ;
- \* the user provides information to the user interface of the parameter selection unit about the desired transformation of  $\beta$  to  $\delta$ ;

- \* the parameter selection unit retrieves R sets of associated data ( $\Sigma_R$ ) from the database, such sets of associated data being selected so that the functionality/functionalities  $^N\beta$  in each set of associated data is/are essentially identical to the functionality/functionalities  $\beta$  in  $^XB$  and the functionality  $^N\delta$  is essentially identical to  $\delta$  in the product  $^XD$ , in order to obtain the R sets of reaction parameters ( $^X\Sigma_R$ ), said R sets of reaction parameters ( $^X\Sigma_R$ ) being accompanied by corresponding information about the chemical substance(s)  $A_R$  under which influence the R reactions should be conducted and information about any additional constituents involved in the chemical reaction;
- \* an array of R reaction mixtures each comprising a predetermined amount of the chemical substance(s)  $A_R$  and the chemical species  $^XB$  and any additional constituents required is prepared according to the sets of reaction parameters;
  - \* each of the R reaction mixtures are treated in the apparatus in accordance with the corresponding set of reaction parameters.
- 2. (Original) A method according to claim 1, wherein the array of R reaction mixtures is provided from  $^{X}B$  stock solution(s) and a kit comprising stock solutions of the chemical substance(s)  $A_{R}$  and any additional constituents required.
- 3. (Original) A method according to claim 1, wherein the R sets of reaction parameters involves the use of more than one chemical substance  $A_R$ .
- 4. (Original) A method according to claim 1, wherein the R sets of reaction parameters involves the use of R chemical substances  $A_R$ .

- 5. (Original) A method according to claim 1, in which the array of R reaction mixtures is prepared by combining the chemical species  ${}^XB$  with the content of one or more of P containers each comprising a chemical substance  $A_R$  including a chemical functionality  $\alpha_R$  which is intended to facilitate the transformation of a functionality  $\beta$  to a functionality  $\delta$  in a chemical reaction involving a chemical species  ${}^XB$ .
- 6. (Original) A method according to claim 1, wherein the R sets of reaction parameters are provided in the form of control parameters for the apparatus.
- 7. (Original) A method according to claim 1, wherein treatment of the R reactions is performed substantially simultaneously.
- 8. (Original) A method according to claim 1, wherein treatment of the R reactions is performed sequentially.
- 9. (Original) A method according to claim 1, wherein the treatment includes heating.
- 10. (Original) A method according to claim 1, wherein the reaction is a microwave facilitated chemical reaction.
- 11. (Original) A method according to claim 1, wherein the apparatus comprises a microwave reaction cavity.

- 12. (Original) A kit for use in the method defined in claim 1, said kit comprising P containers each comprising a chemical substance  $A_R$  including a chemical functionality  $\alpha_R$  which is intended to facilitate the transformation of one or more functionalities  $\beta$  to a functionality  $\delta$  in a chemical reaction involving one or more chemical species  ${}^XB$ , said chemical reaction being intended to result in a reaction product  ${}^XD$  which includes a functionality  $\delta$ , where the chemical reaction involves one or more functionalities  $\beta$  in the  ${}^XB$ 's which are transformed into  $\delta$  in  ${}^XD$ .
- 13. (Original) A kit according to claim 12, which further comprises additional constituents required for the transformation.

## 14-16 (cancelled)

17. (Currently Amended) A kit comprising P containers each comprising a chemical substance  $A_R$  including a chemical functionality  $\alpha_R$  which is intended to facilitate the transformation of one or more functionalities  $\beta$  to a functionality  $\delta$  in a chemical reaction involving one or more chemical species  ${}^XB$ , said chemical reaction being intended to result in a reaction product  ${}^XD$  which includes a functionality  $\delta$ , where the chemical reaction involves one or more functionalities  $\beta$  in the  ${}^XB$ 's which are transformed into  $\delta$  in  ${}^XD$ , said kit usable to conduct R chemical reactions, where R is a positive integer  $\Rightarrow$  1, in a system which includes an apparatus which provides energy for the chemical reactions, said system also including a parameter selecting unit having a user interface and storage means for carrying a database, said chemical reaction involving one or more chemical species  ${}^XB$  and resulting in a reaction product  ${}^XD$  which

includes a functionality  $\delta$ , where the chemical reaction involves one or more functionalities  $\beta$  in the  ${}^XB$ 's which are transformed into  $\delta$  in  ${}^XD$ , each reaction being performed under the influence of one or more corresponding chemical substances  $A_R$ , such chemical substances  $A_R$  including a chemical functionality  $\alpha_R$  being involved in the transformation of the functionality/functionalities  $\beta$  to the functionality  $\delta$ , said database comprising N sets of associated data, each of the N sets comprising

- i) a set of reaction parameters for a chemical reaction involving the transformation of one or more functionalities  ${}^{N}\beta$  of chemical species  ${}^{N}B$  into  ${}^{N}\delta$  in a product  ${}^{N}D$  under the influence of one or more chemical substances  ${}^{N}A$ , such chemical substance(s) each including a chemical functionality  ${}^{N}\alpha$  being involved in the transformation of the functionality  ${}^{N}\beta$  to the functionality  ${}^{N}\delta$ ; and
- ii) functional or structural information about the chemical species <sup>N</sup>B; the method comprising that
- \* the user provides information to the user interface of the parameter selection unit about the functionality/functionalities  $\beta$  in the chemical species  ${}^{X}B$ ;
- \* the user provides information to the user interface of the parameter selection unit about the desired transformation of  $\beta$  to  $\delta$ ;
- \* the parameter selection unit retrieves R sets of associated data ( $\Sigma_R$ ) from the database, such sets of associated data being selected so that the functionality/functionalities  $^N\beta$  in each set of associated data is/are essentially identical to the functionality/functionalities  $\beta$  in  $^XB$  and the functionality  $^{MN}\delta$  is essentially identical to  $\delta$  in the product  $^XD$ , in order to obtain the R sets of reaction parameters ( $^X\Sigma_R$ ), said R sets of reaction parameters ( $^X\Sigma_R$ ) being accompanied by

corresponding information about the chemical substance(s)  $A_R$  under which influence the R reactions should be conducted and information about any additional constituents involved in the chemical reaction;

- \* an array of R reaction mixtures each comprising a predetermined amount of the chemical substance(s) A<sub>R</sub> and the chemical species <sup>X</sup>B and any additional constituents required is prepared according to the sets of reaction parameters;
- \* each of the R reaction mixtures are treated in the apparatus in accordance with the corresponding set of reaction parameters.
- 18. (Previously Presented) A kit according to claim 17, which further comprises additional constituents required for the transformation.
- 19. (Currently Amended) A computer readable data carrier loaded with a computer program system, said computer program system
- \* retrieving information via the user interface of the parameter selection unit about the functionality/functionalities  $\beta$  in the chemical species  ${}^{X}B$ ;
- \* retrieving information via the user interface of the parameter selection unit about the desired transformation of  $\beta$  to  $\delta$ ;
- \* retrieving, via the parameter selection unit, R sets of associated data  $(\Sigma_R)$  from the database, such sets of associated data being selected so that the functionality/functionalities  $^N\beta$  in each set of associated data is/are essentially identical to the functionality/functionalities  $\beta$  in  $^XB$  and the functionality  $^{MN}\delta$  is essentially identical to  $\delta$  in the product  $^XD$ , in order to obtain the R sets of

reaction parameters ( $^{X}\Sigma_{R}$ ), said R sets of reaction parameters ( $^{X}\Sigma_{R}$ ) being accompanied by corresponding information about the chemical substance(s)  $A_{R}$  under which influence the R reactions should be conducted and information about any additional constituents involved in the chemical reaction;

- \* providing instructions to the liquid handler about the preparation of an array of R reaction mixtures each comprising a predetermined amount of the chemical substance(s) A<sub>R</sub> and the chemical species <sup>X</sup>B and any additional constituents required according to the sets of reaction parameters;
- \* providing instructions to the reaction cavity about treatment of each of the R reaction mixtures in the apparatus in accordance with the corresponding set of reaction parameters in order to conduct R chemical reactions, where R is a positive integer  $\rightarrow$  1, in a system which includes an apparatus which provides energy for the chemical reactions, said system also including a parameter selecting unit having a user interface and storage means for carrying a database, said chemical reaction involving one or more chemical species  $^{X}B$  and resulting in a reaction product  $^{X}D$  which includes a functionality  $\delta$ , where the chemical reaction involves one or more functionalities  $\beta$  in the  $^{X}B$ 's which are transformed into  $\delta$  in  $^{X}D$ , each reaction being performed under the influence of one or more corresponding chemical substances  $A_{R}$ , such chemical substances  $A_{R}$  including a chemical functionality  $\alpha_{R}$  being involved in the transformation of the functionality/functionalities  $\beta$  to the functionality  $\delta$ , said database comprising N sets of associated data, each of the N sets comprising
  - i) a set of reaction parameters for a chemical reaction involving the transformation of one or more functionalities  ${}^{N}\beta$  of chemical species  ${}^{N}B$  into  ${}^{N}\delta$  in a product  ${}^{N}D$  under the influence of one or more chemical substances  ${}^{N}A$ , such chemical

substance(s) each including a chemical functionality  $^N\alpha$  being involved in the transformation of the functionality  $^N\beta$  to the functionality  $^N\delta$ ; and

- ii) functional or structural information about the chemical species  ${}^{N}B$ ; the method comprising that
- \* the user provides information to the user interface of the parameter selection unit about the functionality/functionalities  $\beta$  in the chemical species  $^{X}B$ ;
- \* the user provides information to the user interface of the parameter selection unit about the desired transformation of  $\beta$  to  $\delta$ ;
- \* the parameter selection unit retrieves R sets of associated data ( $\Sigma_R$ ) from the database, such sets of associated data being selected so that the functionality/functionalities  $^N\beta$  in each set of associated data is/are essentially identical to the functionality/functionalities  $\beta$  in  $^XB$  and the functionality  $^{MN}\delta$  is essentially identical to  $\delta$  in the product  $^XD$ , in order to obtain the R sets of reaction parameters ( $^X\Sigma_R$ ), said R sets of reaction parameters ( $^X\Sigma_R$ ) being accompanied by corresponding information about the chemical substance(s)  $A_R$  under which influence the R reactions should be conducted and information about any additional constituents involved in the chemical reaction;
- \* an array of R reaction mixtures each comprising a predetermined amount of the chemical substance(s) A<sub>R</sub> and the chemical species <sup>X</sup>B and any additional constituents required is prepared according to the sets of reaction parameters;
  - \* each of the R reaction mixtures are treated in the apparatus in accordance with the corresponding set of reaction parameters.